Soleo – a natural tool to strengthen the plant and it's harvestable produce and increase post harvest quality



David Bernad Technical Manager







Daymsa: presentation

- Daymsa, began manufacturing and marketing operations 31 years old
- Daymsa belongs to a large corporate group called SAMCA.
- ☐ The SAMCA Group is located in Spain, whose activities are focused on the mining, agriculture, energy, plastics, textile and construction sectors. In all, it provides employment for more than 4,000 people. It has facilities all over Spain, as well as in other countries, such as France, Italy or Portugal.
- Daymsa reaches its consumers via a professional distribution network covering Spain and 25 countries of Europe, Africa, Asia and South America, where we have three subsidiaries (Ecuador, Colombia and Brazil).







SOLEO: COMPOSITION

- Plant extract: Mix of natural Rutaceae extracts
- □ Plant strengthener: In Spain under ORDEN APA/1470/2007, otros medios de defensa fitosanitaria.









RECOMMENDED USES (I)

- Protection of the crop against phytopathogenic fungus, specially:
 - Botrytis (Botrytis cinerea)
 - Powdery mildew (e.g. Sphaeroteca macularis, Sphaeroteca fulginea, Leveillula taurica)







RECOMMENDED USES (II)

Extension of post harvest life of fruits and vegetables. Improves the appearance maintaining organoleptic characteristics.











ADVANTAGES (I)

- Soleo can be used as a complement to fungicides
- Alternative to fungicides when they can't be used
 - Harvesting period
 - Organic agriculture
- ☐ Helps to the reduction of pesticide uses
 - **□** Lower number of applications
 - Lower amount of fungicides active substances used on a crop







Main constituents and Mode of action

Natural citric and ascorbic acid, and pectins

- 1. Induces the plant to produce phytoalexins: Immune Response in Plant
- 2. Produces and accumulates antioxidants: Immune Response and Strengthening in plant
- 3. Physical barrier









Application method and doses (I)

Start the treatments as soon as favorable environment condition starts for disease development.

☐ Use foliar application, making sure good wetting of disease susceptible organs.









Application method and doses (II)

- Doses:
 - **▶** Botrytis: 0,5–1,0 cc/l
 - Powdery mildew: 1,0-1,5 cc/l

■ Repeat with a frequency of 10-20 days depending on the environmental conditions







SOLEO: Botrytis cinerea Trial in tomato



Treatments:

- Control (Untreated)
- Soleo 0,5 I/ha
- Soleo 1 l/ha
- Soleo 1,5 l/ha
- Chemical phytosanitary

Location: Andalucía (Spain). Tomato in greenhouse.

Design: Randomized blocks in 4 replicates per treatment. 10 plants/replicate.

a) Applications

First application: When the first signs of disease development. Second application and next: 10-15 days interval.

4 applications were made.

- **b)** Parameters: Incidence and severity in leaves and fruits. Incidence: % of leaves (or fruits) with symptoms in 50 leaves of each repetition. Severity: % of damaged surface in each of 50 leaves (or fruits) of each repetition.
- c) Assessment dates:


Before every application and 14 days after last application.







SOLEO: *Botrytis cinerea* Trial in tomato Results on the leaves



First application: 17/03

Botrytis begin significantly on 09/04


Next applications on assessment dates: 09/04, 21/04 y 02/05







SOLEO: *Botrytis cinerea* Trial in tomato Results on the fruits



First application: 17/03

Botrytis begin significantly on 09/04

Next applications on assessment dates: 09/04, 21/04 y 02/05







Powdery mildew trial in tomato Materials and Methods (1/2)



- Location: Greenhouse in Conil de la Frontera (Cádiz, Spain)
- Cultivar: Tomate var Daniela
- Disease: Leveillula taurica
- Plant density: 25000 p/ha
- Treataments
 - Control (no treated)
 - Soleo 1,0 L/Ha
 - Soleo 1,5 L/Ha
 - Chemical phytosanitary








Materials and Methods (2/2)

- 3 foliar applications:
 - T1: 30/06/2009
 - T2: 14/07/09 = T1+14
 - T3: 30/07/09 = T2+16
- Application equipment
 - The products were sprayed with compressed air sprayers with a lance with cone nozzles and the pressure was 3,5 bar.
 - Spray volume of 1000 L/ha
- Randomized blocks design with 4 replicates
- Each replicate with 14 plants
- Assessments:
 - Incidence and Severity
 - Efficacy (Abbott formula)










































SOLEO: Powdery mildew *Sphaeroteca fulginea* in Cucumber

 Location: Greenhouse in Los Palacios (Sevilla)

Cultivar: Suso

 4 applications. Spray volume in 1st application: 1000 L/Ha. In the 3 successives: 1500 L/Ha.

 Applications: T2=07/05/2008, T3=21/05/2008, T4=04/06/2008

| TREATMENT | DOSAGE |
|---------------------|--------|
| CONTROL (UNTREATED) | |
| SOLEO | 1 L/Ha |
| Phytosanitary | 0.03% |


| PARAMETERS | SAMPLE | DATES |
|---|----------------|---|
| N° LEAVES WITH SYMPTOMS | 20 Leaves/plot | T2 + 0 days T3 + 0 days T4 + 0 days T4 + 10 days |
| % OF ATTACK 0-5 (Towsend & Huberger) | 20 Leaves/plot | T2 + 0 days T3 + 0 days T4 + 0 days T4 + 10 days |
| PHYTOTOXICITY (0-10) | PLOT | T2 + 0 days T3 + 0 days T4 + 0 days T4 + 10 days |








SOLEO: Powdery mildew in Cucumber











SOLEO on lettuce: Botrytis cinerea

- Location: Benicarló (Castellón, ESP)
- Cultivar: Valladolid. Density: 0,3x1,2. Plantation date: 29/12/2009.
- Start of applications: preventive, when the conditions are favorable to development of the disease.
- 4 applications. Spray volume in 1st application:
 500 L/Ha. In 2nd 600 L/ha; in 3^d and 4th: 800 L/Ha.
- Dosage: 0,5 cc/L
- Applications:

| T1+11 | T1+22= T2+11 | T1+32 = T3+10 | T1+40 = T4+8 |
|--------|--------------|---------------|--------------|
| 1/2/10 | 12/2/10 | 22/2/10 | 2/3/10 |










Soleo on Lettuce


Botrytis cinerea:

incidence and

severity results













Soleo trial on Lettuce *Botrytis cinerea:* Damage assessment

Damage assessment (%)

| | Control | Soleo 0,5 cc/l | | |
|---|---------|----------------|----|--|
| 1= No attack | 28 | 61 | 59 | |
| 2= Slight attack, with only the basal stems damaged | 67 | 39 | 41 | |
| 3= No commercial lettuce | 5 | 0 | 0 | |









SOLEO: Powdery mildew in Strawberry

- Objective: check Soleo efficacy on *Sphaeroteca macularis*, in comparison with standard Triadimenol 25% EC.
- Location: Greenhouse in Moguer (Huelva)
- Cultivar: Camarosa
- 4 applications. Spray volume used: 1000
 L/Ha.

| TREATMENTS | DOSAGE |
|---------------------|--------|
| CONTROL (UNTREATED) | • |
| SOLEO | 1 L/Ha |
| TRIADIMENOL 25% EC | 0.05% |









Materials and methods

EC in 'Camarosa' cultivar grown under macrotunnel, in BBCH 89 phenological stage. Plant density was 50.000 plants/ha.

The trial design was set up as a randomized block in 4 replications per treatment (E.P.P.O № 1/104(2)). The row spacing was 1,20 m. The plots had 4 m long, with 2 lines per plot.

The products were sprayed with compressed air sprayers with a lance with cone nozzles and the pressure was 3,5 bar.

Incidence, severity and phytotoxicity were assessed on the following dates: T3+0, T4+0 and T4+10. In previous dates (T1+0, T2+0) not enough disease development was observer in order to make the evaluation.

Powdery mildew incidence was evaluated with the percentaje of affected leaves, assesing 50 leaves per replication. The degree of foliar damage was calculated in accordance with Townsend and Heuberger's formula. For this, severity was evaluated on 50 leaves per replication. Efficacy was calculated using Abbot's formula, based on the degree of damage. Phytotoxicity was assesed with a 0-10 range.


Statistical analysis were assessed using Newman-Keuls Test.







SOLEO: Powdery mildew control in strawberry









SOLEO EFFICACY EVALUATION ON POST-HARVEST DISEASES OF PEACH

Objective: Check out the efficacy of the foliar treatments of the product SOLEO for the control of post-harvest diseases

Trial location: Alcalá del Río (Sevilla, ESP)

Treatments area selection: Peach Sweet Robin cultivar. A band of 15 rows was selected where the different treatments were located.











Treatments:

<u>Treatment A:</u> *Iprodione* (Iprodione 50% w/v CS), 18 days before harvest day (31/03).

<u>Treatment B:</u> A first application of *Iprodione* (Iprodione 50% w/v CS), 18 days before harvest and one application of *SOLEO* on, 2 days before harvest (16/04).

<u>Treatment C:</u> Two applications of SOLEO. A first application 11 days before harvest (7/04) and a second application 2 days before harvest (16/04).

SOLEO dosage: 1 L/ha (1 cc/L).











Podredumbres causadas por patógenos en la muestra de melocotones.

Cladosporium sp.



Cladosporium sp.
Penicillium sp.



Botryosphaeria sp.



Botrytis sp.

Colletotrichum sp.










Post Harvest Trial on peaches: Results after 10 days 20°C (1/2)










Post Harvest Trial on peach: Results 2/2

Tabla 1. Número medio de frutos en porcentaje afectados obtenidos en cada tratamiento. Fecha de evaluación 2/05/08.

| tratamiento | Frutos afectados (%) | | | | | | | | |
|-------------|-------------------------|-----------------|--------------------|-----------------------|-----------------|---------------------|--------------------|-----------------|--------------------|
| | Dañados | Con lesiones | Alternaria s p. | Botryosphaeria sp. | Botrytis sp. | Cladosporium sp. | Colletotrichum sp. | Rhizopus sp. | Penicillium sp. |
| A | 50,0 | 6,0 | 3,0 | 10,2 | 12,0 | 18,0 | 23,0 | 23,0 | 10,0 |
| В | 35,0 | 4,0 | 2,0 | 8,1 | 4,0 | 21,0 | 18,0 | 12,0 | 2,0 |
| С | 47,9 | 1,0 | 12,2 | 9,2 | 9,2 | 34,0 | 25,5 | 14,3 | 1,0 |









Post Harvest Trial on peach: Conclusions

- Soleo applications have permitted obtain a good production of quality fruit erasing the last fungicide
 lprodione treatment and, thus, deleting the risks of pesticides residues in post-harvest period. Only a
 little part of control of Alternaria sp has been reduced, although the Penicillium sp control is widely
 better with two Soleo applications with respect to Iprodione use.
- Soleo treatment made 2 days before harvest did decrease the incidence of all pathogens in evaluated fruits (Alternaria sp, Botryosphaeria sp, Botrytis sp, Colleotrichum sp, Rhizopus sp and Penicillium sp), except for Cladosporium sp, which is the only evaluated pathogen where control has been decreased.
- The replacement of a final Iprodione treatment (18 days before harvest) for 2 Soleo applications, apart from delete the risks of residues for that application, has permitted a similar control of the Botryosphaeria sp, Botrytis sp and Colletotrichum sp post-harvest development.
- The replacement of a final Iprodione treatment (18 days before harvest) for 2 Soleo applications, apart from delete the risks of residues for that application, has reduced the incidency of the Rhizopus sp and Penicilium sp post-harvest development.
- No physiological alterations and no phytotoxicitiy in the crop were observed.
- No spots, derived from product application, were observed on fruits.









Thank you ii





